

It's important to share any impact on your day-to-day activities due to less pain in your feet (eg, time with family, socializing with friends, sleep, cooking, or household chores).

Identify any changes in the treatment notes sections below.

Progress Tracker

Diabetic nerve pain of the feet is different for everyone.

Some people may have pain relief after one treatment, while others may need multiple treatments to feel a difference. Use this progress tracker to help identify your progress over the next few treatments and share it with your healthcare provider.

On the day of each treatment, mark an X on the chart to indicate your pain level on a scale of 0 to 5. Repeat this once a week until your next appointment. It may help to rate your pain on the same day of the week, but pick the day that's right for you.

INDICATION

QUTENZA® (capsaicin) 8% topical system is indicated in adults for the treatment of neuropathic pain associated with postherpetic neuralgia (PHN) and for neuropathic pain associated with diabetic peripheral neuropathy (DPN) of the feet.

IMPORTANT SAFETY INFORMATION

- Treatment with QUTENZA must be performed only by a healthcare provider. You should never apply or remove QUTENZA yourself.

Please see additional Important Safety Information on next page.

Treatment 1 _____

PAIN LEVEL	5	<input type="checkbox"/>												
	4	<input type="checkbox"/>												
	3	<input type="checkbox"/>												
	2	<input type="checkbox"/>												
	1	<input type="checkbox"/>												
	0	<input type="checkbox"/>												
DAY OF TREATMENT	↑		1	2	3	4	5	6	7	8	9	10	11	12
WEEKS AFTER TREATMENT														

Treatment 1 Notes

Treatment 2 _____

PAIN LEVEL	5	<input type="checkbox"/>												
	4	<input type="checkbox"/>												
	3	<input type="checkbox"/>												
	2	<input type="checkbox"/>												
	1	<input type="checkbox"/>												
	0	<input type="checkbox"/>												
DAY OF TREATMENT	↑		1	2	3	4	5	6	7	8	9	10	11	12
WEEKS AFTER TREATMENT														

Treatment 2 Notes

Treatment 3 _____

PAIN LEVEL	5	<input type="checkbox"/>												
	4	<input type="checkbox"/>												
	3	<input type="checkbox"/>												
	2	<input type="checkbox"/>												
	1	<input type="checkbox"/>												
	0	<input type="checkbox"/>												
DAY OF TREATMENT	↑		1	2	3	4	5	6	7	8	9	10	11	12
WEEKS AFTER TREATMENT														

Treatment 3 Notes

Pain levels: 0 = no pain at all and 5 = most pain imaginable.

INDICATION

QUTENZA[®] (capsaicin) 8% topical system is indicated in adults for the treatment of neuropathic pain associated with postherpetic neuralgia (PHN) and for neuropathic pain associated with diabetic peripheral neuropathy (DPN) of the feet.

IMPORTANT SAFETY INFORMATION

- Treatment with QUTENZA must be performed only by a healthcare provider. You should never apply or remove QUTENZA yourself.
- Do not touch QUTENZA or items exposed to capsaicin. Touching QUTENZA and then accidentally touching other areas of your body can cause severe irritation of eyes, mucous membranes, respiratory tract, and skin.
- Do not touch your eyes or other unintended target areas during the QUTENZA application. QUTENZA is not for use near eyes or mucous membranes. Do not sniff or inhale near QUTENZA as this may cause you to cough or sneeze. If irritation of eyes or airways occurs or any side effects become severe, notify your doctor immediately.
- The treated area may be sensitive to heat (e.g., hot showers/bath, direct sunlight, vigorous exercise) for a few days following treatment.
- You may experience substantial pain during the treatment. Tell your healthcare provider if you are experiencing pain; a cool compress or medicine for the pain can be provided to help lessen your discomfort.
- QUTENZA can cause serious side effects, including pain and increases in blood pressure during or right after treatment. Your healthcare provider should check your blood pressure during treatment with QUTENZA. If you have high blood pressure that is not well controlled by medicine, or have had recent heart problems, stroke, or other vascular problems, you may be at increased risk and should discuss with your doctor whether QUTENZA is right for you.
- Tell your doctor if you have reduced sensation in the feet. You may notice that you have less feeling for hot or sharp pain where QUTENZA was applied, but this is usually minor and temporary.

The most common side effects of QUTENZA are redness, pain, or itching where QUTENZA was applied. You should tell your doctor if any side effects bother you or do not go away.

To report suspected adverse reactions, contact Averitas Pharma, Inc. at 1-877-900-6479 (Option 1) or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

For more information, ask your healthcare provider or pharmacist.

Please see full [Prescribing Information](#).